




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07C 229/42, A61K 31/215 C07C 237/20, A61K 31/16	A1	(11) International Publication Number: WO 94/04484 (43) International Publication Date: 3 March 1994 (03.03.94)
(21) International Application Number: PCT/EP93/01906 (22) International Filing Date: 20 July 1993 (20.07.93) (30) Priority data: MI92A002006 20 August 1992 (20.08.92) IT (71) Applicants (for all designated States except US): CORLAY S.L. [ES/ES]; Calle Azcona, 31, E-28028 Madrid (ES). METGROVE LTD. [IE/IE]; Earlsfort Centre, Earlsfort Terrace, 17, Dublin 2 (IE). (72) Inventors; and (75) Inventors/Applicants (for US only) : MATJI, José, Antonio [ES/ES]; ALCAIDE, Antonio [ES/ES]; Corlay S.L., Calle Azcona, 31, E-28028 Madrid (ES).		(74) Agent: TRUPIANO, Roberto; Brevetti Europa s.r.l., Piazza Bernini, 6, I-20133 Milano (IT). (81) Designated States: BR, CA, JP, KR, RU, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: NITRIC ESTERS OF DERIVATIVES OF THE 2-(2,6-DI-HALOPHENYLAMINO)PHENYLACETIC ACID AND PROCESS FOR THEIR PREPARATION

**(57) Abstract**

Object of this invention are nitric esters of derivatives of the 2-(2,6-di-halophenylamino)phenylacetic acid, having general formula (I), as well as their pharmaceutical utilization and process for their preparation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

NITRIC ESTERS OF DERIVATIVES OF THE 2-(2,6-DI- HALO-PHENYLAMINO)PHENYLACETIC ACID AND PROCESS FOR THEIR PREPARATION

TECHNICAL FIELD

Object of the present invention are nitric esters of derivatives of 2-(2,6-di-halo-phenylamino)phenylacetic acid, their pharmaceutical utilization and process for their preparation.

PRIOR ART

The sodium salt of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid has been used for a long time in the pharmaceutical field for its anti-inflammatory activity and has been sold throughout the world for many years. The process for its preparation has been described in the Dutch Patent application No. 6.604.752 and in the US Patent No. 3.558690.

The pharmacological profile and the effectiveness of the sodium salt of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid are described in Am.J.Med.80, Suppl. 4B, 1-87 (1986), while other data concerning its pharmacological activity as anti-inflammatory agent are reported, for instance, in C.A.74, 86215 m (1971); Krupp et al. Experimentia 29,450 (1973).

The utilization of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid as an anti-inflammatory preparation causes, as known, very severe adverse reactions, for instance in the gastro-intestinal apparatus, as well as damages to the liver and the kidneys. There exist numerous experimental evidences [S.MONCADA,

R.M.J.PALMER, E.A.HIGGS, Pharmacological Reviews, 43(2), 109-142 (1991); T.F.LUSHER, C.M.BOULANGER, Y.DOHI, Z.YANG, Hypertension, 19, 117-130 (1992)], on whose basis the integrity of the vasal endothelium is
5 assumed to act as a basically important protective barrier to prevent the onset of pathologic reactions in various organs and apparatuses.

Such protective barrier, and therefore the integrity of the vasal endothelium, is ensured, on the physiological
10 plane, by the presence of nitric oxide and prostacyclin.

The treatment with drugs having an anti-inflammatory activity, such as, for instance, the sodium salt of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid, causes
15 the inhibition of the cyclo-oxygenase, an enzyme which governs the synthesis of the prostacyclin precursor.

As a consequence, the production of prostacyclin being in this way inhibited, the tissular reserve of same is markedly depauperated, with ensuing compromission of
20 the vasal endothelium.

As said, because of this endothelial damage due to the reduction of prostacyclin, diffuse pathologic reactions break out which affect the gastrointestinal apparatus, the kidneys and the liver.

25

OBJECTS OF THE INVENTION

Object of the present invention is to provide a product which, while ensuring the maintenance of the


pharmacological activity characteristic of the known anti-inflammatory preparations, can also eliminate the adverse reactions caused by the treatment with said drugs.

- 5 Another object of the present invention is the realization of a process for the preparation of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid having an anti-inflammatory activity and that are exempt from those adverse reactions that are typical of
10 the anti-inflammatory drugs.

DESCRIPTION OF THE INVENTION

These and still other objects and advantages which shall appear from the following description are obtained by derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid, which derivatives,
15 according to the present invention, have the following general formula:

20



25

wherein:

A and B are selected among hydrogen, linear or branched,

substituted or non substituted alkyl chains, X is a halogen selected among chloride and bromine, Y is selected among oxygen, NH, NR_1 , wherein R_1 is a linear or branched alkyl group and n is comprised between 1 and 10.

5 In fact, it has been observed that the introduction of a group such as a terminal nitric ester in the derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid as in (I) permits to preserve the pharmacological activity of anti-inflammatory drugs, while eliminating the
10 adverse reactions caused by the treatment with said drugs.


It has also been noticed that the derivatives (I) are useful for the treatment of different unhealthy conditions, such as for instance rheumatic diseases in
15 general, immunologic disorders, and that they can also alleviate painful conditions of low-middle severity of any kind.

Besides, the derivatives (I) subject matter of the present invention are useful in the treatment of the
20 illnesses of the cardiovascular apparatus and in particular in the treatment of myocardial and brain ischaemiae, as well as in cases of arterial thrombosis.

Always according to the present invention, a nitric ester of a derivative of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I) proved to be especially advantageous, wherein:
25


A and B are hydrogen, X is chlorine, Y is oxygen, and n

is equal to four, according to the following formula:



10 Also particularly advantageous according to this invention is a nitric ester of a derivative of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I), wherein:


A and B are hydrogen, X is chlorine, Y is oxygen, and n
15 is equal to two, according to the following formula:



25 For the preparation of the derivatives (I) of the 2-(2,6-di-halo-phenylamino)phenylacetic acid subject matter of this invention, a first process proved to be particularly advantageous which, according to the

present invention, comprises the following phases:

- Reaction between the sodium salt of the 2-(2,6-di-halo-phenylamino)phenylacetic acid or of the 2-(2,6-di-halo-phenylamino)phenylacetic acid functionalized to the carboxylic group, and a compound having the following general formula:




wherein:

R_4 is selected among chlorine, bromine, NHR in which R is hydrogen or linear or branched alkyl chain, A and B are selected among hydrogen, linear or branched, substituted or non substituted alkyl chains, R_3 is selected among chlorine, bromine and iodine; and n is comprised between 1 and 10, the carboxylic group of the 2-(2,6-di-halo-phenylamino)phenylacetic acid being functionalized as acyclic chloride, anhydride or the like, obtaining in this way the corresponding monomer ester or the corresponding amide;

- Reaction of said monomer ester or of said corresponding amide with a nitrating agent such as AgNO_3 or the like, obtaining in this way nitric esters or derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I).

A second process proved also particularly advantageous which, always according to the present invention, comprises the following phases:

- Reaction between the sodium salt of the 2-(2,6-di-halo-phenylamino)phenylacetic acid or of the 2-(2,6-di-halo-phenylamino)phenylacetic acid functionalized to the carboxylic group, with a compound having the following general formula:



wherein:

R_4 is selected among chlorine, bromine, NHR in which R is hydrogen or linear or branched alkyl chains, A and B are selected among hydrogen, linear or branched, substituted or non substituted alkyl chains, and n is comprised between 1 and 10, the carboxylic group of the 2-(2,6-di-halo-phenylamino)phenylacetic acid being functionalized as acyclic chloride, anhydride or the like, obtaining in this way either the corresponding monomer ester or the corresponding amide;

- Reaction of said monomer ester or said corresponding amide with an halogenating compound such as PBr_3 or the like, obtaining in this way said monomer ester or said amide, characterized by the presence of a terminal


halogen group;

- Reaction of said monomer ester or said amide, characterized by the presence of a terminal halogen group with a nitrating agent such as AgNO_3 or the like, obtaining in this way nitric esters of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I).


The solvents which are utilized in the processes subject matter of this invention are preferably selected among chloroform, methylene chloride, acetonitrile, dimethylformamide, tetrahydrofuran, dioxan and the like.

Such processes for the preparation of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I), subject matter of the present invention, consist of a limited number of phases, allowing to obtain the products deriving from such processes rapidly, with satisfactory yields and to high amounts, even on an industrial basis.

According to the processes subject matter of this invention, the preparation of nitric esters of derivatives of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid proved particularly advantageous, having the following formulae:



and



which are prepared as described in the following examples, which are given as mere indications that do not limit in any way the protection scope of this invention.

EXAMPLE 1

a) 6 g of 1-Br-4-Cl-butane diluted in 250 ml of dimethylformamide were dripped to a solution of 10 g of sodium salt of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid in 100 ml of dimethylformamide. The reaction mix was stirred for 12 hours at room temperature, then diluted with water and extracted with methylene chloride. The so extracted organic phase was anhydried on sodium sulfate and the solvent was low-

pressure evaporated until 14 g of dry residual were obtained.

The residual was purified by chromatography on silica gel, utilizing chloroform as eluant system.

5 The head fractions were then collected, and by low-pressure evaporation of the solvent 11 g of dry residual were obtained and then chromatographed anew on silica gel, utilizing an eluant mix constituted by hexane/ether 7/3 (v/v).

10 The head fractions were collected, the solvent was low-pressure evaporated, and 3 g of 2-(2,6-di-chloro-phenylamino)phenylacetate of 4-chlorobutyl (VIII) were obtained.

IR (cm^{-1}): C=O, 1741; NH, 3340.

15 ^1H -NMR(300MHz) (CDCl_3): 1.9 ppm(m, 4H); 3.6 ppm(m, 2H);
3.85 ppm(s, 2H); 4.2 ppm(m, 2H); 6.5-7.45 ppm(m, aromatics).

Mass spectrometry (i.e) : M^+ 385

20 b) 1.2 g of AgNO_3 diluted in 11 ml of acetonitrile were dripped to 2 g of (VIII) obtained as described in a), diluted in 7 ml of acetonitrile. The reaction mix was stirred for 12 hours at the temperature of 85°C and then filtered.

25 The solvent was low-pressure evaporated from the resulting solution, and a residual was obtained to which 30 ml of methylene chloride were added. The mix so obtained

ned was filtered anew, the organic phase was water-washed and then anydried on sodium sulfate. The solvent was low-pressure evaporated and 2.8 g of dry residual were obtained, which were purified thereafter by chromatography on silica gel, utilizing an eluant mix constituted by hexane/ether 7/3 (v/v). The fractions containing the product were collected, the solvent was low-pressure evaporated and 2.5 g of nitric ester of 2-(2,6-di-chloro-phenylamino)phenylacetate of 4-hydroxibutyl (II) were obtained.

IR (cm^{-1}): C=O, 1729; NH, 3322; ONO_2 , 1637.

^1H -NMR(80 MHz) (CDCl_3): 1.75 ppm (m, 4H); 3.8 ppm (s, 2H); 4.2 ppm (m, 2H); 4.4 ppm (m, 2H); 6.45-7.4 ppm (m, aromatics).

Mass spectrometry (i.e.) M^+ 412.

EXAMPLE 2

a) 0,5 g of ethylester of the 2-(2,6-di-chlorophenylamino)phenylacetic acid were added to 0,5 ml of 4-aminobutanol and the mix so obtained was stirred at the temperature of 100°C for 12 hours. The mix was then brought again to room temperature, diluted with 5 ml of water and extracted with 5 ml of methylen chloride. The organic phase so extracted was anydried on sodium sulfate and the solvent was low-pressure evaporated until 0,19 g of 2-(2,6-di-chloro-phenylamino)-4-hydroxibutyl-phenylacetamide (XII) were obtained.

IR (cm^{-1}) (nujol): C=O, 1648; NH and OH, 3413.

$^1\text{H-NMR}$ (80 MHz) (CDCl_3): 1.65 ppm (m, 4H); 3.3 ppm (m, 2H); 3.6 ppm (m, 2H); 6.08 ppm (m, 1H); 6.5 ppm (dd, 1H); 6.85-7.5 ppm (m, 6H).

Mass spectrometry: PM 366

5 b) 1,14 g of PBr_3 were added to a solution of 0,19 g of 2-(2,6-di-chloro-phenylamino)-4-hydroxybutyl-phenylacetamide (XII) in 10 ml of chloroform; the mix so obtained was stirred for 30 minutes and then diluted with 10 ml of water. The organic phase was separated
10 and anhydried on sodium sulfate, and then the solvent was low-pressure evaporated, obtaining in this way a raw residual which was purified by chromatography, utilizing an eluant mix constituted by methylene chloride/ethyl acetate 10/0,1 (v/v).

15 The intermediate fractions were recovered, the solvent was low-pressure evaporated and 50 mg of 2-(2,6-di-chloro-phenylamino)-4-bromobutyl-phenylacetamide (XIII) were obtained.

$^1\text{H-NMR}$ (80 MHz) (CDCl_3): 1.73 ppm (m, 4H); 3.3 ppm (m, 8H); 3.67 ppm (s, 2H); 5.91 ppm (broad s, 1H); 6.5 ppm (dd, 1H); 6.92-7.29 ppm (m, 5H); 7.4 ppm (d, 1H).
20

c) 1.5 g of AgNO_3 diluted in 10,7 ml of acetonitrile were added to a solution constituted by 2,8 g of 2-(2,6-di-chloro-phenylamino)-4-bromobutyl-phenylacetamide (XIII) diluted in 9 ml of acetonitrile. The reaction
25 mix was stirred at the temperature of 25°C for 3 days and then filtered. The solvent was low-pressure evapo-


rated from the resulting solution, obtaining in this way a residual which was purified by chromatography, utilizing methylene chloride as eluant. The fractions containing the product were collected, the solvent was low-pressure evaporated, and 0,5 g of nitric ester of 2-(2,6-di-chloro-phenylamino)-4-hydroxybutylphenylacetamide (XI) were obtained.


IR(cm^{-1}) (nujol): C=O, 1650; NH, 3290; ONO_2 , 1630.

^1H -NMR (80 MHz) (CDCl_3): 1.62 ppm (m, 4H); 3.28 ppm (m, 2H); 4.4 ppm (t, 2H); 5.3 ppm (broad s, 1H); 6.49 ppm (dd, 1H); 6.85-7.36 ppm (m, 5H); 7.4 ppm (d, 1H).

Mass spectrometry: PM 411.

There has been determined by means of biologic tests, the anti-inflammatory and analgesic activity for instance of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I) having the following formulae:





10 The anti-inflammatory activity of said derivatives of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid has been determined in Wistar rats, by utilizing the method of the carrageenan edema, as reported in C.A.WINTER, E.RISLEY, G.W.NUSS, Proc. Soc. Exp.Biol.Med. 111, 544-547 (1962), while the analgesic activity of said derivatives has been determined in Swiss mice, as reported by 15 L.C.HENDERSHOT, J.FORSAITH, J.Pharmacol.Exp.Ter. 125, 237-249 (1959).

20 The anti-inflammatory and analgesic activity of said derivatives is given on Table 1, and is expressed as a power ratio relative to 2-(2,6-di-chloro-phenylamino)phenylacetic acid taken as a reference.

Each value represents a mean of the values obtained by the treatment of 10 animals.

25 The compounds (II) and (XI) utilized for said biological tests were suspended in 0.5% carboxymethylcellulose before the administration.

TABLE 1

COMPOUND	ANTI-INFLAMM. ACTIVITY	ANALGESIC ACTIVITY
----------	------------------------	--------------------

XI	1.25	1.40
II	1.30	1.50
2-(2,6-di-chloro-phenylamino)phenyl-		
acetic acid	1	1

Then, the acute toxicity of said derivatives (II) and (XI) was evaluated by oral administration of a single dose of each compound (II) and (XI), utilizing for each derivative groups of 10 Swiss mice. The lethality incidence and the onset of toxic symptoms were assessed within a period of observation of 14 days.

Even upon administration of 250 mg/kg of the compound (II) or the compound (XI) no apparent toxicity symptoms have been observed in the studied animals.

Further biological experiments, suitable to determine the pharmacotoxicologic profile of the derivative (II) have been carried out by examining said derivative (II) in comparison with the 2-(2,6-di-chloro-phenylamino)phenylacetic acid taken as a reference.

A. PHARMACODYNAMIC ACTIVITY

Acute models

RAT CARRAGEENAN PAW EDEMA: the values of ED 30 (mg/kg p.o.) obtained are respectively equal to 4,88 for the compound (II) and to 4,21 for the 2-(2,6-di-chloro-phenylamino)phenylacetic acid, showing a comparable effectiveness between the two compounds.

MOUSE PHENYLQUINONE WRITHING: at doses ranging between

3 and 10 mg/kg p.o., the derivative (II) has shown a full effectiveness and its potency resulted almost comparable to that of the 2-(2,6-di-chloro-phenylamino)phenylacetic acid and of indomethacin.

5 Subacute models

RAT ADJUVANT ARTHRITIS: the animals treated for 19 consecutive days (from the 3rd to the 21th day after the adjuvant injection) with 3,0 mg/kg p.o. of 2-(2,6-di-chloro-phenylamino)phenylacetic acid or with 1,5 or
10 3,0 mg/kg p.o. of the compound (III), have shown a significant reduction of the arthritic symptomatology.

Rat gastrointestinal tolerability

In all the animals treated with 15 mg/kg p.o. of 2-(2,6-di-chloro-phenylamino)phenylacetic acid, severe
15 diffuse ulcerations have been observed; small ulcers have been observed also in animals treated with 3,5 and 7,0 mg/kg p.o.

The average dose of ulcerogenicity for the 2-(2,6-di-chloro-phenylamino)phenylacetic acid has been calculated as being equal to 6,1 mg/kg p.o.
20

The compound (II) showed to be very well tolerated even at much higher doses compared to the above mentioned ones; small ulcers have been noticed only in 2 animals out of 10, treated with 100 mg/kg. Therefore
25 it was impossible to determine the average ulcerogenicity dose for the compound (II).

General pharmacology

Secondary pharmacological evaluations of the compound (II) have been carried out by comparison with the 2-(2,6-di-chloro-phenylamino)phenylacetic acid; no additional effects have been observed besides the primary pharmacological activity on the central nervous system, the autonomic system, the cardiovascular, respiratory and gastrointestinal systems.

B. TOXICOLOGY

Acute toxicity in rodents

Studies have been carried out in two animal species and following two different administration routes.

The following values of LD50 (mg/kg) and of the 95% fiducial limits have been obtained:

rat, oral route: 511 (356-732); mouse, oral route: 497 (323-762); rat, intraperitoneal route: 237 (156-359); mouse, intraperitoneal route: 253 (171-374).

Maximum tolerated dose in non-rodents

The compound (II) was very well tolerated in this animal species which, as known, is particularly sensitive to this class of compounds.

The animals have been treated with doses increasing from 250 to 1000 mg/kg of compound (II): the lowest dose caused no symptomatology, the intermediate dose caused only a reversible diarrhea, while the highest dose caused a severe but reversible diarrhea. On the contrary, the administration in the same conditions of 10 mg/kg of 2-(2,6-di-chloro-phenylamino)phenylacetic

acid caused the death of the animals.

Subacute toxicity in rodents

5 The animals have been treated with 5, 15 and 30 mg/kg of compound (II) for 4 weeks. The general conditions and clinical behaviour, body weight gain, water and food consumption, hematology and clinical chemistry have shown that the two lowest doses have been well tolerated.

Subacute toxicity in the dog


10 The animals have been treated with 5, 15 and 30 mg/kg of compound (II) for 4 weeks. The general conditions and clinical behaviour, body weight gain, water and food consumption, hematology and clinical chemistry have shown that the two lowest doses have been well
15 tolerated.

20

25

CLAIMS

1. Derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid, characterized in that they have the following general formula:



wherein:

A and B are selected among hydrogen, linear or branched, substituted or non substituted alkyl chains, X is an halogen selected among chlorine and bromine, Y is selected among oxygen, NH, NR₁, wherein R₁ is a linear or branched alkyl group, and n is comprised between 1 and 10.

2. A derivative of the 2-(2,6-di-halo-phenylamino)phenylacetic acid according to claim 1, characterized in that X is chlorine, A and B are hydrogen, Y is oxygen and n is equal to 4.

3. A derivative of the 2-(2,6-di-halo-phenylamino)phenylacetic acid according to claim 1, characterized in that X is chlorine, A and B are hydrogen, Y is oxygen and n is equal to 2.

4. A derivative of the 2-(2,6-di-halo-phenylamino)phenylacetic acid according to claim 1,


characterized in that X is chlorine, A and B are hydrogen, Y is NH and n is equal to 4.

5. Derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid according to claim 1, characterized in that they are utilizable in the pharmaceutical field as anti-inflammatory drugs.

6. Derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid according to claim 1, characterized in that they are utilizable in the treatment of rheumatic diseases, in the treatment of immunological disorders and of middle severity painful conditions.

7. Derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid according to claim 1, characterized in that they are utilizable in the treatment of illnesses of the cardiovascular apparatus, of myocardial an brain ischaemiae and in the arterial thrombosis.


8. A process for the preparation of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid, having the following general formula:



wherein:

A and B are selected among hydrogen, linear and branched, substituted and non substituted alkyl chains, X is an halogen selected among chlorine and bromine, Y is selected among oxygen, NH, NR₁, wherein R₁ is a linear or branched alkyl group and n is comprised between 1 and 10, characterized in that it comprises the following phases:

- Reaction between the sodium salt of the 2-(2,6-dihalo-phenylamino)phenylacetic acid or of the 2-(2,6-dihalo-phenylamino)phenylacetic acid functionalized to the carboxylic group, with a compound having the following general formula:




wherein:

R₄ is selected among chlorine, bromine, NHR in which R is hydrogen or linear or branched alkyl chain, A and B are selected among hydrogen, linear or branched, substituted or non substituted alkyl chains, R₃ is selected among chlorine, bromine and iodine, and n is comprised between 1 and 10, the carboxylic group of the 2-(2,6-dihalo-phenylamino)phenylacetic acid being functionalized as acilic chloride, anhydride or the like, obtaining in this way the corresponding monomer ester or the corre-

sponding amide;

- Reaction of said monomer ester or said amide with a nitrating agent such as AgNO_3 or the like, obtaining in this way nitric esters of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid (I).

9. Process for the preparation of derivatives of the 2-(2,6-di-halo-phenylamino)phenylacetic acid, having the following general formula:




wherein:

A and B are selected among hydrogen, linear or branched, substituted or non substituted alkyl chains, X is an halogen selected among chlorine and bromine, Y is selected among oxygen, NH, NR₁, wherein R₁ is a linear or branched alkyl group and n is comprised between 1 and 10, characterized in that it comprises the following phases:

- Reaction between the sodium salt of the 2-(2,6-di-halo-phenylamino)phenylacetic acid or of the 2-(2,6-di-halo-phenylamino)phenylacetic acid functionalized to the carboxylic group, with a compound having the

following general formula:



5

wherein:

R_4 is selected among chlorine, bromine, NHR in which R is hydrogen or linear or branched alkyl chain, A and B are
10 selected among hydrogen, linear or branched, substituted or non substituted alkyl chains, and n is comprised between 1 and 10, the carboxylic group of the 2-(2,6-di-halo-phenylamino)phenylacetic acid being functionalized as acilic chloride, anydride or the
15 like, obtaining in this way the corresponding monomer ester or the corresponding amide;


- Reaction of said monomer ester or said amide with an halogenating compound such as PBr_3 or the like, obtaining in this way said monomer ester or said amide,
20 characterized by the presence of a terminal halogen group;

- Reaction of said monomer ester or said amide, characterized by the presence of a terminal halogen group with a nitrating agent such as AgNO_3 or the like,
25 obtaining in this way nitric esters of derivatives of the 2-(2,6-di-halo-phenylamine)phenylacetic acid (I).

NITRIC ESTERS OF DERIVATIVES OF THE 2-(2,6-DI-HALO-PHENYLAMINO)PHENYLACETIC ACID AND PROCESS FOR THEIR PREPARATION

ABSTRACT

Object of this invention are nitric esters of derivatives of the 2-(2,6-di-halophenylamino)phenylacetic acid, having the following general formula:



as well as their pharmaceutical utilization and process for their preparation.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07C229/42 A61K31/215 C07C237/20 A61K31/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,3 558 690 (A. SALLMANN ET AL.) 26 January 1971 cited in the application see claims ---	1-9
A	WO,A,91 06539 (AMERICAN HOME PRODUCTS CORPORATION) 16 May 1991 see claims ---	1-9
A	DE,A,34 07 507 (A. NATTERMANN & CIE GMBH) 5 September 1985 see claims -----	1-9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

28 October 1993

Date of mailing of the international search report

- 9. 11. 93

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

SANCHEZ Y GARCIA, J

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-3558690	26-01-71	US-A- 3895063	15-07-75
		US-A- 3804877	16-04-74
		BE-A- 696248	29-09-67
		CH-A- 473769	15-06-69
		CH-A- 526512	15-08-72
		DE-A,B,C 1618465	27-05-71
		FR-M- 6680	03-02-69
		FR-A- 1517251	
		GB-A- 1132318	
		NL-C- 137301	
		NL-A- 6704484	02-10-67
		BE-A- 679315	10-10-66
		CH-A- 460804	
		DE-A- 1793592	20-04-72
		DE-A- 1543639	05-02-70
		FR-M- 5524	06-11-67
		FR-A- 1487352	
		GB-A- 1139332	
		NL-C- 133740	
		NL-A- 6604752	10-10-66
		BE-A- 725792	20-06-69
		CH-A- 487116	15-03-70
		CH-A- 487841	31-03-70
		DE-A- 1815807	24-07-69
		FR-M- 8235	28-09-70
		NL-A- 6817964	24-06-69
		BE-A- 725793	20-06-69
		CH-A- 487840	31-03-70
		DE-A- 1815802	10-07-69
		FR-A- 1595382	08-06-70
		GB-A- 1257190	15-12-71
		NL-A- 6817965	24-06-69
		SE-B- 366296	22-04-74
		CH-A- 485667	15-02-70
WO-A-9106539	16-05-91	US-A- 5021576	04-06-91
		AU-B- 640429	26-08-93
		AU-A- 6724790	31-05-91
		CA-A- 2067135	28-04-91
		EP-A- 0523046	20-01-93

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/EP 93/01906

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9106539		JP-T- 5502022 US-A- 5159085	15-04-93 27-10-92
DE-A-3407507	05-09-85	NONE	

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)